DOI: 10.5281/zenodo.17634590

CHAPTER 5

From Soil to Solutions: The Role of Actinomycetes in Interdisciplinary Life Science Research

Meenakshi S.,[1] Jyothi Hiremath,[2] Shivaveerakumar S.[1]*

[1]Department of Microbiology, Davangere University, Shivagangothri, Davanagere, Karnataka, India

[2]Department of Food Science and Nutrition, Khaja Banda Nawaz University, Kalaburagi, Karnataka, India

Corresponding author Email: shiv_math1984@yahoo.com

Received: 10 October 2025; Accepted: 27 October 2025; Available online: 17 November 2025

Abstract: Actinomycetes are filamentous bacteria celebrated for their extraordinary secondary-metabolite diversity, accounting for most naturally derived antibiotics and numerous anticancer, antiviral, and immunomodulatory compounds. Yet their significance transcends pharmacology. Through complex ecological interactions and metabolic versatility, actinomycetes participate in nutrient cycling, bioremediation, plant growth promotion, and biomaterial synthesis. This chapter traces the scientific journey of actinomycetes from their soil origins to their contemporary roles in interdisciplinary life-science research highlighting how omics technologies, synthetic biology, and

This work is licensed under a Creative Commons Attribution 4.0 International License. This allows re-distribution and re-use of a licensed work on the condition that the author is appropriately credited and the original work is properly cited.

Interdisciplinary Research in Life Sciences: A Path Towards Sustainability (Vol. 3) - Jayvardhan V. Balkhande & Jalander Vaghmare (Eds.)

ISBN: 978-93-95369-86-2 (paperback) 978-93-95369-58-9 (electronic) | © 2025 Advent Publishing.

https://doi.org/10.5281/zenodo.17634590

environmental engineering are reshaping our capacity to harness these microorganisms for sustainable solutions.

Keywords: Streptomyces, bio-active molecules, Bio-synthetic gene clusters, secondary metaboloites.

1. Introduction

The soil ecosystem hosts an immense microbial diversity, estimated at more than 10^4 bacterial species per gram.[1] Within this microscopic metropolis, actinomycetes stand out as filamentous saprophytes that produce earthy aromas (geosmin) and a staggering array of bioactive molecules. Their discovery redefined medicine: the isolation of streptomycin from *Streptomyces griseus* by Selman Waksman in 1943 inaugurated the antibiotic era.[2]

Over subsequent decades, more than two-thirds of all naturally derived antibiotics have been traced to actinomycetes, particularly the genus *Streptomyces.* However, the story of these microbes is no longer confined to drug discovery. Modern genomics reveals that actinomycetes encode thousands of "silent" or cryptic biosynthetic gene clusters (BGCs) whose products remain unexplored.[3] Advances in metagenomics, transcriptomics, and metabolomics are awakening this hidden potential, making actinomycetes indispensable to interdisciplinary science linking molecular genetics, chemical ecology, environmental sustainability, and bioengineering.[4]

2. Taxonomy and Morphological Diversity

Actinomycetes belong predominantly to the order Actinomycetales within the class Actinobacteria. They exhibit high genomic G + C content (60–75 %), filamentous growth, and spore formation reminiscent of fungi, though they are true prokaryotes. The major genera include *Streptomyces, Nocardia, Micromonospora, Frankia, Rhodococcus, Actinomyces*, and *Saccharopolyspora.*[5]

Morphologically, *Streptomyces* species form extensively branched substrate and aerial mycelia that differentiate into spores under nutrient limitation, a process controlled by the bld (bald) and whi (white) regulatory genes. The complexity of this developmental cycle mirrors that of filamentous fungi, highlighting an evolutionary convergence in soil colonization strategies.[6]

Recent phylogenomic analyses employing 16S rRNA sequencing and average nucleotide identity (ANI) clustering have refined actinomycete taxonomy, revealing deep evolutionary splits between terrestrial and marine clades. High-resolution comparative genomics also suggests frequent horizontal gene transfer events, enabling the acquisition of biosynthetic pathways across species. This genetic fluidity underpins their chemical creativity and environmental adaptability.[7]

2.1. Evolutionary Insights

Molecular-clock studies indicate that actinomycetes diverged early in bacterial evolution, likely over 1 billion years ago, contemporaneous with the colonization of terrestrial habitats by early fungi[8]. Fossilized actinobacterial filaments have been detected in Precambrian cherts, suggesting that these microbes contributed to the first soil formation events.

Their evolutionary success arises from modular polyketide synthase (PKS) and non-ribosomal peptide synthetase (NRPS) systems, which function as natural combinatorial chemistries.[9] The shuffling and recombination of these modules across lineages have generated unparalleled metabolic diversity. Moreover, regulatory genes such as *afsR, adpA*, and *sigB* coordinate secondary-metabolite biosynthesis with stress responses, enabling actinomycetes to adapt rapidly to environmental fluctuations.[10&11]

3. Ecology and Environmental Adaptations

Actinomycetes are ubiquitous, colonizing soils, sediments, compost, freshwater, and marine habitats. Their ecological functions are multifaceted:

3.1. Decomposition and Nutrient Cycling

They secrete extracellular enzymes cellulases, chitinases, xylanases, and lignin peroxidases that decompose complex polymers of plant and animal origin.[10] Through mineralization of organic matter, they recycle carbon, nitrogen, and phosphorus, sustaining soil fertility.

3.2. Antagonism and Chemical Signaling

In competitive soil microhabitats, actinomycetes produce antibiotics and volatile organic compounds (VOCs) that suppress rival microbes.[12] Some VOCs, such as geosmin and 2-methylisoborneol, act as signaling molecules influencing microbial succession and even attracting soil invertebrates that aid spore dispersal.[6]

3.3. Symbioses and Mutualisms

The genus Frankia forms nitrogen-fixing nodules with over 200 species of non-leguminous plants, including *Casuarina* and *Alnus*.[10] Endophytic *Streptomyces* species inhabit root tissues, promoting plant growth by secreting indole-3-acetic acid (IAA), siderophores, and antifungal metabolites.[13]

3.4. Stress Tolerance and Extremophily

Actinomycetes from deserts, glaciers, and marine sediments withstand salinity, UV radiation, and desiccation through protective pigments (melanin, carotenoids) and osmolytes (trehalose,

ectoine).[14&15] Thermophilic genera such as *Thermobifida* and *Thermomonospora* produce thermostable enzymes crucial for industrial catalysis.[16]

3.5. Ecological Resilience and Carbon Sequestration

Recent metagenomic studies indicate that actinomycetes contribute significantly to long-term soil carbon stabilization by producing recalcitrant microbial necromass that binds to mineral particles.[17] This positions them as essential agents in climate-resilient ecosystems.[12]

4. Metabolomic Diversity and Secondary Metabolite Biosynthesis

Actinomycetes are often termed the "chemical factories" of nature. Their genomes harbor between 20 and 50 biosynthetic gene clusters (BGCs), encoding diverse secondary metabolites including antibiotics, antitumor agents, antivirals, immunosuppressants, pigments, and enzymes.[18]

4.1. Types of Metabolites and Pathways

The biosynthetic machinery of actinomycetes is dominated by polyketide synthases (PKSs) and non-ribosomal peptide synthetases (NRPSs) giant, modular enzymatic complexes that assemble complex molecules from simple acyl-CoA and amino acid precursors.[19&20]

Polyketides like erythromycin and avermectin arise from sequential condensation of acetate and propionate units, while NRPS products such as actinomycin and vancomycin are peptide-based but synthesized independently of ribosomes.[21]

Hybrid PKS–NRPS systems (e.g., bleomycin) further expand chemical diversity through domain recombination.[4]

4.2. Genetic Regulation and Cryptic Pathways

Actinomycete genomes contain numerous "silent" BGCs that are not expressed under laboratory conditions. Genome mining tools such as antiSMASH and PRISM predict the existence of hundreds of cryptic clusters.[22] New strategies—like co-culture induction, epigenetic modulation, and heterologous expression—are now routinely used to activate these pathways.[23]

Transcriptional regulators (*afsR, adpA*) and global signaling molecules like γ-butyrolactones coordinate the transition between primary metabolism and secondary metabolism.[24] The integration of metabolomics with transcriptomics has helped map entire metabolic networks, offering blueprints for synthetic biology.[25]

4.3. Natural Product Libraries and Chemical Space

As of current date, more than 30,000 actinomycete metabolites have been structurally characterized.[10] These molecules cover unique chemical scaffolds, often with stereochemical complexity beyond synthetic chemistry's reach.[6] Their structural novelty continues to feed drug-discovery pipelines worldwide.

5. Medical Biotechnology and Pharmaceutical Applications

5.1 Antibiotics: The Cornerstone of Modern Medicine

The first medically significant antibiotic, streptomycin, was discovered from *Streptomyces griseus* in 1943.[26] Since then, *Streptomyces* and related genera have yielded tetracycline, chloramphenicol, erythromycin, vancomycin, rifamycin, and daptomycin, among many others.[27]

Actinomycete-derived antibiotics target all major bacterial pathways protein synthesis, cell-wall formation, nucleic-acid replication, and membrane integrity.[28] Their continued relevance persists even in the era of synthetic drugs because resistance mechanisms are constantly evolving in pathogens.[1]

Recent bioprospecting of marine actinomycetes, particularly *Salinispora tropica* and *Micromonospora marina*, has uncovered new antibiotic classes such as salinosporamide A, a potent proteasome inhibitor used in cancer therapy.[29] Genome editing with CRISPR-Cas9 and recombineering now allows tailoring antibiotic scaffolds for improved activity and reduced toxicity.[30]

5.2 Anticancer and Immunomodulatory Compounds

The contribution of actinomycetes to oncology is equally profound. *Streptomyces peucetius* produces doxorubicin and daunorubicin, which intercalate DNA and inhibit topoisomerase II.[31] *Streptomyces hygroscopicus* gave us rapamycin, an mTOR inhibitor that revolutionized immunosuppressive therapy and cancer biology.[9]

Metabolomic profiling has revealed that actinomycetes produce numerous cytotoxic macrolides, depsipeptides, and enediynes active against multidrug-resistant cancer cell lines.[32]

5.3 Antiviral and Antifungal Agents

In recent years, actinomycetes have produced novel antiviral compounds active against influenza, SARS-CoV-2, and HIV replication. Polycyclic macrolactams and alkaloids derived from Streptomyces species inhibit viral polymerases or interfere with host entry.[22]

Similarly, actinomycete antifungal metabolites; like natamycin, nystatin, and candicidin remain mainstays in treating fungal keratitis and candidiasis.[20]

5.4 Overcoming Antimicrobial Resistance (AMR)

With antimicrobial resistance emerging as a global crisis, actinomycetes are pivotal in providing next-generation antimicrobials.[26] Combinatorial biosynthesis and synthetic biology now enable the generation of novel analogues (e.g., modified glycopeptides) that evade bacterial resistance mechanisms. Machine-learning-guided BGC mining is also identifying new leads in silico before lab validation.[20]

6. Agricultural Biotechnology and Plant Health

Actinomycetes are integral to sustainable agriculture, promoting plant growth, enhancing nutrient uptake, and suppressing pathogens.

6.1. Biofertilization and Phytohormones

Many *Streptomyces* species secrete plant growth–promoting hormones such as indole-3-acetic acid (IAA), gibberellins, and cytokinins, stimulating root elongation and seed germination.[21]

6.2. Biocontrol of Plant Pathogens

Actinomycetes produce antifungal antibiotics (e.g., amphotericin, actinomycin) and hydrolytic enzymes that degrade fungal cell walls. *Streptomyces lydicus* and *S. avermitilis* are commercial biocontrol strains against *Rhizoctonia, Fusarium*, and *Pythium*.[2]

6.3. Siderophore and VOC Production

Siderophores such as desferrioxamine chelate iron in the rhizosphere, depriving pathogens while supporting plant nutrition. Meanwhile, volatile organic compounds (2-pentylfuran, geosmin) act as interkingdom signals that enhance systemic resistance in plants.[33]

6.4. Soil Health and Microbiome Engineering

Actinomycetes stabilize soil aggregates and release extracellular polysaccharides that improve structure and water retention. Synthetic ecology experiments have shown that engineered consortia of actinomycetes can suppress root pathogens while promoting beneficial mycorrhizae.[34]

Table 1: Representative Actinomycetes and Their Agricultural Applications

Genus	Key Metabolite	Function	Reference
Streptomyces lydicus	Natamycin	Antifungal biocontrol	[35]
Frankia alni	Nitrogenase complex	N_2 fixation in non-legumes	[36]
Micromonospora aurantiaca	IAA	Root elongation	[37]
Rhodococcus erythropolis	Biosurfactants	Soil remediation and nutrient uptake	[38]

7. Industrial Enzyme Production and Bioprocessing

Actinomycetes are prolific producers of extracellular enzymes: cellulases, proteases, lipases, amylases, chitinases, and xylanases used in textiles, detergents, food processing, and biofuels.[39]

Modern strain engineering (adaptive evolution, CRISPR editing) has increased enzyme yield and thermostability.[34]

Streptomyces rimosus produces a highly stable alkaline protease used in detergent industries, while *Thermobifida fusca* secretes thermostable cellulases crucial for lignocellulosic bioethanol production. Enzyme immobilization on nanocarriers further improves reusability in industrial reactors.[40]

7.1. Industrial Bioprocess Optimization

Optimization of fermentation conditions carbon/nitrogen ratio, pH, and aeration is essential for maximizing yields. Systems biology tools now enable metabolic flux analysis and predictive control of actinomycete bioreactors.[41]

8. Environmental Biotechnology and Bioremediation

Actinomycetes play critical roles in environmental detoxification. They degrade hydrocarbons, pesticides, heavy metals, and xenobiotics through versatile catabolic pathways.

8.1. Hydrocarbon Degradation

Rhodococcus species metabolize long-chain alkanes, aromatic hydrocarbons, and even crude oil via oxygenase-mediated oxidation.[42] They are essential in oil-spill bioremediation and wastewater treatment.

8.2. Pesticide and Plastic Biodegradation

Streptomyces viridosporus produces lignin peroxidases capable of degrading organophosphates and polyethylene films. Recent work demonstrates enzymatic breakdown of polylactic acid (PLA) plastics by thermophilic actinomycetes, presenting sustainable waste solutions.[6,19&]

8.3. Heavy-Metal Biotransformation

Actinomycetes can bioaccumulate or transform metals like cadmium, lead, and chromium through chelation and redox reactions. Biosorption studies using *Nocardia* biomass have achieved up to 90 % removal efficiency of cadmium from contaminated effluents.[43]

8.4. Microbial Fuel and Carbon Capture Interfaces

Electroactive actinomycetes produce conductive pigments (melanin) that facilitate electron transfer in microbial fuel cells.[44] Their role in soil carbon sequestration further aligns with climate-resilient biotechnologies.[45]

9. Omics Revolution in Actinomycete Research

The advent of high-throughput sequencing and omics technologies has revolutionized actinomycete biology, revealing a genomic and metabolic landscape of staggering complexity.[23]

9.1. Genomics

Whole-genome sequencing of *Streptomyces coelicolor* and *S. avermitilis* revealed that over 5–8% of their genomes are devoted to secondary metabolism. Comparative genomics now guides the discovery of new biosynthetic gene clusters (BGCs) by identifying conserved modular motifs.[46]

Advances in long-read sequencing have facilitated assembly of complex actinomycete genomes, often exceeding 9 Mbp. Genome mining tools such as antiSMASH, MIBiG, and ClusterFinder enable rapid annotation of cryptic BGCs.[47]

9.2. Transcriptomics

RNA-seq analyses provide snapshots of gene expression under different environmental stimuli or co-culture conditions. They have shown that many "silent" clusters are transcriptionally activated only under nutrient stress or interspecies competition. Integration of transcriptomic and metabolomic data now allows precise correlation between gene expression and metabolite production.[48]

9.3. Proteomics

Proteomic profiling using LC-MS/MS and iTRAQ quantitation identifies key enzymes involved in precursor biosynthesis and pathway regulation.[7] Such studies have revealed global regulatory proteins like AfsR, PhoP, and BldD as master switches for secondary metabolism.[49&50]

9.4. Metabolomics

NMR-based and MS-based metabolomics map the chemical fingerprints of actinomycete cultures. Coupled with stable isotope labeling, metabolomics elucidates fluxes through central and secondary metabolic pathways. This systems-level insight enables metabolic engineering strategies for maximizing bioactive compound yield[51].




Figure 1: "Integration of multi-omics approaches in *Streptomyces* metabolism." [25]

10. Synthetic Biology and Systems Biotechnology

Synthetic biology has ushered in a new era of rational manipulation of actinomycete genomes to enhance metabolite discovery and production.

10.1. Gene Cluster Refactoring

Traditional random mutagenesis is being replaced by precise refactoring of biosynthetic clusters. Heterologous expression in optimized chassis strains like *Streptomyces coelicolor* M145 or *Saccharopolyspora erythraea* allows activation of silent BGCs.[52] Modular cloning systems such as Golden Gate and CRISPR-assisted recombineering streamline this process.[37&52]

10.2. Combinatorial Biosynthesis

Mixing and matching PKS and NRPS modules from different actinomycetes has generated "unnatural natural products" with new pharmacological properties. The production of hybrid macrolides and glycopeptides through domain swapping exemplifies how biosynthetic engineering can outperform natural evolution.[53]

10.3. Metabolic Engineering

Systems-level manipulation of precursor supply pathways such as malonyl-CoA and methylmalonyl-CoA pools significantly boosts secondary metabolite yields. Flux balance analysis and genome-scale metabolic models (GEMs) now allow prediction and optimization of metabolic bottlenecks.[54&55]

10.4. Genome Editing and Regulation

CRISPR-Cas9, base editors, and CRISPRi systems have been customized for actinomycetes, enabling precise gene knock-ins, knockouts, and repression of competing pathways. Synthetic promoters and tunable riboswitches further fine-tune expression dynamics, creating programmable biosynthetic systems.[56]




Figure 2: "Schematic view of CRISPR-mediated activation of silent gene clusters in *Streptomyces.*"[56]

11. Actinomycetes in Nanotechnology and Bioelectronics

Actinomycetes' biomineralization capabilities and redox-active metabolites have inspired advances in nanotechnology, biosensors, and bioelectronics.[57]

11.1. Biogenic Nanoparticle Synthesis

Several *Streptomyces* and *Rhodococcus* species reduce metal ions to nanoparticles; silver, gold, selenium, and zinc oxide through enzymatic and extracellular polymeric matrix reactions. These biogenic nanoparticles exhibit potent antimicrobial, antioxidant, and catalytic activities.[58]

Streptomyces sp. NH1 and *Streptomyces viridogens* HM10 for example, synthesizes silver nanoparticles under controlled conditions, which have been used as antimicrobial and antibacterial agents respectively.[59&57] The marine Streptomyces synthesized AgNps demonstrated notable acaricidal activity against *Rhipicephalus microplus* and *Haemaphysalis bispinosa.*[60]

11.2. Conductive Biomolecules and Bioelectronics

Melanin and phenazine pigments produced by certain actinomycetes act as natural semiconductors, facilitating electron transfer. Such materials are being integrated into microbial fuel cells and bioelectronic sensors. The coupling of these biomolecules to electrode surfaces improves current density and energy recovery.[61]

11.3. Biosensors and Biointerfaces

Actinomycete enzymes such as cholesterol oxidase and peroxidases serve as biosensing elements for detecting glucose, cholesterol, and pollutants. The robustness and thermostability of these enzymes make them ideal for industrial sensor development.[62]

Table 2: Actinomycete-Derived Nanomaterials and Applications

Source	Nanomaterial	Application	Reference
Streptomyces sp. NH1	Ag nanoparticles	Antimicrobial activity	57
Streptomyces viridogens HM10	Ag nanoparticles	Antibacterial	59
Streptomyces sp. LK3	Ag nanoparticles	Acaricidal activity	60
Rhodococcus sp.	Ag nanoparticles	Antimicrobial coatings	58

12. Actinomycetes' ecological functions extend into climate resilience, soil carbon dynamics, and green biotechnologies.

12.1. Carbon Cycling and Sequestration

By decomposing lignocellulosic residues and producing humic precursors, actinomycetes contribute to stable soil organic matter formation. Their extracellular polysaccharides bind soil particles, enhancing aggregate stability and reducing CO_2 emissions.[63]

12.2. Methane and Nitrogen Cycling

Recent metagenomic surveys have identified actinobacterial lineages harboring methane monooxygenase-like genes, suggesting potential roles in methane oxidation. Similarly, *Frankia* species contribute to biological nitrogen fixation in non-leguminous trees, critical for reforestation and climate mitigation.[64]

12.3. Climate-Smart Agriculture

Bioinoculants containing *Streptomyces* and *Micromonospora* improve drought tolerance and nutrient uptake in crops like wheat and rice under climate stress. These formulations enhance root biomass and photosynthetic efficiency, reducing fertilizer dependency.[65]

12.4. Bioplastic and Green Chemistry

Actinomycetes synthesize polyhydroxyalkanoates (PHAs) and extracellular polysaccharides as biodegradable plastics. Industrial-scale production using *Rhodococcus jostii* has achieved yields comparable to petrochemical plastics. The enzymes they produce also catalyze green synthesis of fine chemicals, replacing hazardous reagents.[66]

13. Interdisciplinary Collaborations in Life Sciences

The study of actinomycetes sits at the intersection of multiple disciplines: microbiology, chemistry, genomics, environmental science, and bioengineering.

13.1. Bioinformatics and Data Science

Integration of omics datasets through machine learning enables predictive discovery of new metabolites. AI assisted BGC clustering has already led to identification of over 200 novel compounds in silico. [67]

13.2. Chemical Ecology and Systems Design

Actinomycete VOCs and antibiotics influence entire soil microbial networks. Interdisciplinary research combining metabolomics, soil ecology, and plant physiology is redefining our understanding of rhizosphere dynamics.[68]

13.3. Biomedical Engineering

Cross-disciplinary collaborations between microbiologists and material scientists have resulted in the use of actinomycete-derived polymers and nanoparticles for wound dressings, biosensors, and scaffolds.[69]

13.4. Environmental and Space Microbiology

NASA-funded studies have shown that certain Streptomyces strains survive simulated Mars conditions, making them potential models for astrobiology and closed-loop bioregenerative life support systems.[70]

14. Future Prospects and Emerging Frontiers

The trajectory of actinomycete research is rapidly expanding from traditional microbiology to encompass cutting-edge interdisciplinary technologies. The next decade will likely witness a fusion of AI-driven discovery, synthetic biology, and eco-systems engineering.[71]

14.1. Artificial Intelligence and Predictive Genomics

Machine-learning algorithms are revolutionizing the prediction of biosynthetic gene clusters (BGCs) and the classification of secondary metabolites. Deep-learning models trained on large genomic datasets can now infer the structure and potential bioactivity of uncharacterized metabolites directly from sequence data. AI-based dereplication tools are accelerating the identification of novel compounds, reducing redundancy in natural product discovery.[72]

14.2. Metagenomics and Cultivation of Rare Actinomycetes

The majority of actinomycetes in nature remain uncultivable. Novel culturomics techniques diffusion chambers, soil substrate membrane systems, and microfluidic isolation are unlocking this "microbial dark matter." These approaches have led to the recovery of rare genera such as *Catenulispora*, *Actinopolymorpha*, and *Planctonospora* with unique metabolic traits.[73]

14.3. Genome Editing and Modular Pathway Design

The coupling of CRISPR systems with machine-guided pathway design enables *de novo* assembly of modular biosynthetic pathways. Such synthetic platforms could generate custom libraries of therapeutic molecules and industrial enzymes, transforming drug discovery into a programmable science.[54]

14.4. Actinomycetes and Human Microbiome Research

Emerging evidence indicates that commensal actinobacteria (e.g., *Cutibacterium*, *Corynebacterium*) play beneficial roles in the skin and gut microbiomes. Understanding their interactions with host immunity may yield next-generation probiotics and microbiome-based therapies.[74]

14.5. Global Sustainability and Climate Adaptation

Actinomycetes hold vast potential for circular bioeconomy models. By converting agricultural residues into bioplastics, degrading environmental pollutants, and sequestering soil carbon, they align perfectly with United Nations Sustainable Development Goals (SDGs). International collaborations linking microbiologists, environmental engineers, and policy scientists are essential to scale these innovations.[6]

Conclusion

From their humble beginnings as soil decomposers to their current stature as molecular innovators, actinomycetes represent the very essence of interdisciplinary biology. Their metabolic ingenuity, ecological adaptability, and evolutionary resilience make them indispensable allies in addressing the grand challenges of the 21st century antibiotic resistance, food security, environmental degradation, and climate change.

Modern genomics and synthetic biology are unlocking a new generation of actinomycete-derived solutions: programmable antibiotics, sustainable biomaterials, and green industrial processes. Yet, this progress must be guided by ecological awareness and responsible biotechnology.

In essence, actinomycetes exemplify a simple truth: the solutions to some of humanity's most complex problems are often buried in the soil beneath our feet.

References

1 S. Meenakshi, J. Hiremath, M. H. Meenakshi and S. Shivaveerakumar, *J. Pure Appl. Microbiol.*, 2024, **18**, 118–143.

2 S. A. Waksman, H. C. Reilly and D. A. Harris, *J. Bacteriol.*, 1948, **56**, 259–269.

3 N. T. Quach, T. H. Nguyen Vu, T. L. Bui, A. T. Pham, T. T. An Nguyen, T. T. Xuan Le, T. T. Thuy Ta, P. Dudhagara and Q.-T. Phi, *Pol. J. Microbiol.*, 2022, **71**, 381–394.

4 D. A. Hopwood, *Streptomyces in nature and medicine: the antibiotic makers*, Oxford University Press, Oxford ; New York, 2007.

5 S. Shivaveerakumar, M. Dn, H. Raghavendra and A. Dayanand, *J. Pure Appl. Microbiol.*, 2014, 8.

6 R. V. Rai and J. A. Bai, Eds., *Natural Products from Actinomycetes: Diversity, Ecology and Drug Discovery*, Springer Singapore, Singapore, 2022.

7 H.-P. Fiedler, *Nat. Prod. Lett.*, 1993, **2**, 119–128.

8 M. Dworkin, S. Falkow, E. Rosenberg, K.-H. Schleifer and E. Stackebrandt, Eds., *The Prokaryotes*, Springer New York, New York, NY, 2006.

9 E. Stegmann and Y. Mast, Eds., *Actinomycetes: The Antibiotics Producers*, MDPI - Multidisciplinary Digital Publishing Institute, Basel, Switzerland, 2020.

10 E. A. Barka, P. Vatsa, L. Sanchez, N. Gaveau-Vaillant, C. Jacquard, H.-P. Klenk, C. Clément, Y. Ouhdouch and G. P. van Wezel, *Microbiol. Mol. Biol. Rev.*, 2015, **80**, 1–43.

11 X. Huang, D. Pinto, G. Fritz and T. Mascher, *J. Bacteriol.*, 2015, **197**, 2517–2535.

12 O. Tyc, C. Song, J. S. Dickschat, M. Vos and P. Garbeva, *Trends Microbiol.*, 2017, **25**, 280–292.

13 T. T. Nguyen, T. T. Nguyen, T. H. Nguyen, L. T. Nguyen, D. T. Tran, S. T. Dinh, T. M. Vu and C. X. Nguyen, *Sci. Rep.*, 2025, **15**, 7049.

14 K. Ndabankulu, S. O. Egbewale, Z. Tsvuura and A. Magadlela, *Sci. Rep.*, 2022, **12**, 12601.

15 P. M. Plonka and M. Grabacka, *Acta Biochim. Pol.*, 2006, **53**, 429–443.

16 J. Seabury, J. Salvaggio, J. Domer, J. Fink and T. Kawai, *J. Allergy Clin. Immunol.*, 1973, **51**, 161–173.

17 G. Xu, X. Kang, F. Wang, W. Zhuang, W. Yan and K. Zhang, *Commun. Earth Environ.*, 2024, **5**, 397.

18 R. Asah-Asante, L. Tang, X. Gong, S. Fan, C. Yan, J. O. Asante and Q. Zeng, *World J. Microbiol. Biotechnol.*, 2025, **41**, 211.

19 J. Wink, F. Mohammadipanah and J. Hamedi, Eds., *Biology and Biotechnology of Actinobacteria*, Springer International Publishing, Cham, 2017.

20 K. Alam, A. Mazumder, S. Sikdar, Y.-M. Zhao, J. Hao, C. Song, Y. Wang, R. Sarkar, S. Islam, Y. Zhang and A. Li, *Front. Microbiol.*, DOI:10.3389/fmicb.2022.968053.

21 C. Tringali, Ed., *Bioactive compounds from natural sources: isolation, characterisation, and biological properties*, Taylor & Francis, London New York, 2001.

22 D. Dhanasekaran and Y. Jiang, Eds., *Actinobacteria - Basics and Biotechnological Applications*, InTech, 2016.

23 N. Lee, S. Hwang, Y. Lee, S. Cho, B. Palsson and B.-K. Cho, 2019, **29**, 667–686.

24 Q. Fang, F. Maglangit, L. Wu, R. Ebel, K. Kyeremeh, J. H. Andersen, F. Annang, G. Pérez-Moreno, F. Reyes and H. Deng, *Molecules*, 2020, **25**, 460.

25 Hindra, P. Pak and M. A. Elliot, *J. Bacteriol.*, 2010, **192**, 4973–4982.

26 R. E. De Lima Procópio, I. R. Da Silva, M. K. Martins, J. L. De Azevedo and J. M. De Araújo, *Braz. J. Infect. Dis.*, 2012, **16**, 466–471.

27 J. K. Schniete and L. T. Fernández-Martínez, *Current Opinion in Microbiology*, 2024, **79**, 102487.

28 S. N, D. D, S. N and S. A, *Virol. Immunol. J.*, 2025, **9**, 1–8.

29 S. K. Sr and K. V. B. Rao, *Asian Pac. J. Trop. Biomed.*, 2012, **2**, 787–792.

30 L. Karthik, Ed., *Actinobacteria: Microbiology to Synthetic Biology*, Springer Nature Singapore, Singapore, 2022.

31 J. S. Yaradoddi, M. H. Kontro and S. V. Ganachari, Eds., *Actinobacteria: Ecology, Diversity, Classification and Extensive Applications*, Springer Nature Singapore, Singapore, 2021.

32 L. Ryan, B. Lamarre, T. Diu, J. Ravi, P. J. Judge, A. Temple, M. Carr, E. Cerasoli, B. Su, H. F. Jenkinson, G. Martyna, J. Crain, A. Watts and M. G. Ryadnov, *J. Biol. Chem.*, 2013, **288**, 20162–20172.

33 T. Arai and Y. Mikami, *Appl. Microbiol.*, 1972, **23**, 402–406.

34 S. Siddharth, R. R. Vittal, J. Wink and M. Steinert, *Microorganisms*, 2020, **8**, 225.

35 H. Wu, W. Liu, L. Shi, K. Si, T. Liu, D. Dong, T. Zhang, J. Zhao, D. Liu, Z. Tian, Y. Yue, H. Zhang, B. Xuelian and Y. Liang, *Sci. Rep.*, 2017, **7**, 9114.

36 I. Gifford, S. Vance, G. Nguyen and A. M. Berry, *Front. Microbiol.*, 2019, **10**, 2230.

37 A. Hirsch and M. Valdes, *Soil Biol. Biochem.*, 2010, **42**, 536–542.

38 G. J. Pacheco, E. M. P. Ciapina, E. de B. Gomes and N. P. Junior, *Braz. J. Microbiol.*, 2010, **41**, 685–693.

39 D. Majidi and N. Aksöz, *Am. J. Microbiol. Res.*, 2013, **1**, 1–3.

40 S. Shivaveerakumar and D. Agsar, *Indian J. Biotechnol.*, **2021**, *20*, 47.

41 M. Rudrappa, S. Kumar M, R. S. Kumar, A. I. Almansour, K. Perumal and S. Nayaka, *Microbiol. Res.*, 2022, **263**, 127130.

42 F. Peng, Z. Liu, L. Wang and Z. Shao, *J. Appl. Microbiol.*, 2007, **102**, 1603–1611.

43 L. Karthik, *Protocols of Actinomycetes: Microbiology to Gene editing*, CRC Press, Boca Raton, 1st edn., 2024.

44 M. Al Khatib, M. Harir, J. Costa, M. C. Baratto, I. Schiavo, L. Trabalzini, S. Pollini, G. M. Rossolini, R. Basosi and R. Pogni, *Molecules*, 2018, **23**, 1916.

45 N. E.-A. El-Naggar and W. I. A. Saber, *Polymers*, 2022, **14**, 1339.

46 J. K. Jansson and K. S. Hofmockel, *Curr. Opin. Microbiol.*, 2018, **43**, 162–168.

47 P. Szychowiak, K. Villageois-Tran, J. Patrier, J.-F. Timsit and É. Ruppé, *Ann. Intensive Care*, 2022, **12**, 3.

48 Z. Liu, Y. Zhao, C. Huang and Y. Luo, *Front. Bioeng. Biotechnol.*, 2021, **9**, 632230.

49 A. Manteca, J. Sanchez, H. R. Jung, V. Schwämmle and O. N. Jensen, *Mol. Cell. Proteomics MCP*, 2010, **9**, 1423–1436.

50 Y. Wang, L. Zhang, X. Zhao, H. Li and D. H. Sherman, *PLoS Biol.*, **2024**, *22*, e3002528.

51 B. Wang, F. Guo, S.-H. Dong and H. Zhao, *Nat. Chem. Biol.*, 2018, **15**, 111.

52 The Sound of Silence: Activating Silent Biosynthetic Gene Clusters in Marine Microorganisms, https://www.mdpi.com/1660-3397/13/8/4754?utm_source=chatgpt.com, (accessed 9 October 2025).

53 D. G. Kim, B. Gu, Y. Cha, J. Ha, Y. Lee, G. Kim, B.-K. Cho and M.-K. Oh, *Nat. Commun.*, 2025, **16**, 874.

54 J.-X. Ma, W.-Y. He, H.-M. Hua, Q. Zhu, G.-S. Zheng, A. A. Zimin, W.-F. Wang and Y.-H. Lu, *ACS Synth. Biol.*, 2023, **12**, 3114–3123.

55 Y. Wu, H. Jin, Q. Yu, Z. Wei, J. Zhu, X. Qiu, G. Luo, J. Li, Y. Zhan, D. Cai and S. Chen, *Appl. Environ. Microbiol.*, 2025, **91**, e01953-24.

56 M. M. Zhang, F. T. Wong, Y. Wang, S. Luo, Y. H. Lim, E. Heng, W. L. Yeo, R. E. Cobb, B. Enghiad, E. L. Ang and H. Zhao, *Nat. Chem. Biol.*, 2017, **13**, 607–609.

57 M. Heinemann, C. Rosa, G. Rosa and D. Dias, *Trends Environ. Anal. Chem.*, 2021, **30**, e00129.

58 M. S. Kuyukina, M. V. Makarova, I. B. Ivshina, K. P. Kazymov and B. M. Osovetsky, *Int. J. Mol. Sci.*, 2022, **23**, 12939.

59 R. Balagurunathan, M. Radhakrishnan, R. B. Rajendran and D. Velmurugan, *Indian J. Biochem. Biophys.*, 2011, **48**, 331–335.

60 K. Loganathan, G. Kumar, V. K. Arivarasan, A. N. Rajeshkumar and B. Rao, *Bioprocess Biosyst. Eng.*, **2013**, *37*, 209.

61 T. Arai and Y. Mikami, *Appl. Microbiol.*, 1972, **23**, 402–406.

62 M. G. Hassan, M. T. Hawwa, D. M. Baraka, H. M. El-Shora and A. A. Hamed, *BMC Microbiol.*, 2024, **24**, 21.

63 A. R. G. Mason, M. J. Salomon, A. J. Lowe and T. R. Cavagnaro, *J. Clean. Prod.*, 2023, **417**, 137993.

64 S. Kim Tiam, H. Boubakri, L. Bethencourt, D. Abrouk, P. Fournier and A. Herrera-Belaroussi, *Genes*, 2023, **14**, 530.

65 B. Benmrid, C. Ghoulam, Y. Zeroual, L. Kouisni and A. Bargaz, *Commun. Biol.*, 2023, **6**, 1016.

66 R. S. K. Sachan, A. Kumar, A. Karnwal, P. Paramasivam, A. Agrawal and A. G. Ayanie, *BMC Microbiol.*, 2025, **25**, 136.

67 K. L. Howe, K. W. Seitz, L. G. Campbell, B. J. Baker, J. C. Thrash, N. N. Rabalais, M.-K. Rogener, S. B. Joye and O. U. Mason, *FEMS Microbiol. Ecol.*, 2022, **99**, fiac153.

68 L. Cuervo, C. Méndez, J. A. Salas, C. Olano and M. G. Malmierca, *Cells*, 2022, **11**, 3510.

69 M. Vinosha, S. Palanisamy, R. Muthukrishnan, S. Selvam, E. Kannapiran, S. You and N. M. Prabhu, *Process Biochem.*, 2019, **85**, 219–229.

70 NASA Astrobiology, https://astrobiology.nasa.gov/news/microbes-could-survive-thin-air-of-mars/?utm_source=chatgpt.com, (accessed 9 October 2025).

71 R. C. Lynch, S. P. Smith and J. B. Hanson, *ISME J.*, **2014**, *8*, 1.

72 M. Cortesão, K. Siems, S. Koch, K. Beblo-Vranesevic, E. Rabbow, T. Berger, M. Lane, L. James, P. Johnson, S. M. Waters, S. D. Verma, D. J. Smith and R. Moeller, *Front. Microbiol.*, DOI:10.3389/fmicb.2021.601713.

73 K. L. Howe, K. W. Seitz, L. G. Campbell, B. J. Baker, J. C. Thrash, N. N. Rabalais, M.-K. Rogener, S. B. Joye and O. U. Mason, *FEMS Microbiol. Ecol.*, 2022, **99**, fiac153.

74 S. M. J. Mortazavi, I. Said-Salman, A. R. Mortazavi, S. El Khatib and L. Sihver, *Front. Microbiol.*, 2024, **14**, 1237564.